Table 4. Rank and % composition of key forage species in ungulate diets, Jarbidge Mountain, Nevada, 1999.

Key Species	Elk		Elk		Deer		Sheep		Cattle	
	Spring		Summer		Summer		Summer		Summer	
	Rank	%	Rank	%	Rank	%	Rank	%	Rank	%
Arrowleaf balsamroot	36	0.3	19	0.8	18	1.3	0	0.0	0	0.0
Bluebunch wheatgrass	2	7.6	5	3.5	0	0.0	3	11.8	1	20.3
Curlleaf mountain mahogany	4	7.2	3	10.0	28	0.4	0	0.0	0	0.0
Idaho fescue	17	1.5	0	0.0	0	0.0	6	4.3	8	4.0
Kentucky bluegrass	0	0.0	0	0.0	0	0.0	5	8.3	4	9.4
Lupine spp.	5	5.5	1	32.0	1	31.9	2	13.6	14	2.2
Mountain brome	9	3.4	12	1.8	31	0.3	12	2.5	15	2.1
Mountain snowberry	20	1.5	0	0.0	0	0.0	0	0.0	0	0.0
Needlegrass spp.	18	1.5	6	3.5	23	0.6	4	11.7	2	12.7
Sandberg's bluegrass	3	7.2	4	3.7	26	0.5	1	16.0	3	10.9
Snowbrush ceanothus	1	21.6	2	16.3	2	11.6	21	0.8	0	0.0
% Total Diet		57.0		71.6		46.6		69.0		61.6